

Purification, crystallization and preliminary X-ray analysis of two nudix hydrolases from *Deinococcus radiodurans*

Holbrook *et al.*

Synopsis

Two putative nudix hydrolases, DR1025 and DR0079, from the radioresistant bacterium *D. radiodurans* have been expressed, purified, crystallized and their diffraction data collected to 1.4 and 1.9 Å resolution, respectively. Anomalous diffraction data were measured to 1.8 Å resolution for a Sm³⁺ derivative of DR1025.

Keywords: nudix hydrolase; *Deinococcus radiodurans*; DNA maintenance.

Queries and comments

Please supply or correct as appropriate all **bold underlined** text.

Subject index

The following terms will be used to index your paper. Authors wishing to recommend additional index entries should give these below.

Hydrolases
nudix hydrolases (*D. radiodurans*)

Author index

Authors' names will normally be arranged alphabetically under their family name and this is commonly their last name. Prefixes (van, de *etc.*) will only be taken into account in the alphabetization if they begin with a capital letter. Authors wishing their names to be alphabetized differently should indicate this below.

Holbrook, E.L.
Schulze-Gahmen, U.
Buchko, G.W.
Ni, S.
Ni, S.
Kennedy, M.A.
Kennedy, M.A.
Holbrook, S.R.

Purification, crystallization and preliminary X-ray analysis of two nudix hydrolases from *Deinococcus radiodurans*

Elizabeth L. Holbrook,^{a,b} Ursula Schulze-Gahmen,^a Garry W. Buchko,^c Shuisong Ni,^c Michael A. Kennedy^c and Stephen R. Holbrook^{a*}

^aStructural Biology Department, Physical Biosciences Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA, ^bChemistry Department, University of California at Berkeley, Berkeley, CA 94720, USA, and ^cFundamental Sciences, Biology Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99352, USA

Correspondence e-mail: srholbrook@lbl.gov

Received 30 October 2002

Accepted 30 January 2003

Two nudix hydrolases from *Deinococcus radiodurans* have been purified and crystallized. Diffraction data have been collected to 1.4 and 1.9 Å resolution for DR1025 and DR0079, respectively. DR1025 belongs to space group $P4_12_12/P4_32_12$, with unit-cell parameters $a = b = 53.2$, $c = 122.6$ Å (unit-cell volume 346 883 Å³, $V_M = 2.5$ Å³ Da⁻¹, solvent content 50.2%). DR0079 belongs to space group $C222_1$, with unit-cell parameters $a = 34.1$, $b = 157.2$, $c = 126.5$ Å (unit-cell volume 677 308 Å³, $V_M = 2.2$ Å³ Da⁻¹, solvent content 44.0%). The calculated cell content of DR1025 indicates the presence of one molecule in the asymmetric unit. Dynamic light scattering suggests it to be a monomer in solution. The space group and unit-cell parameters of DR0079 indicate the presence of two molecules per asymmetric unit. Gel filtration and NMR spectroscopy suggest it to be a monomer in solution.

1. Introduction

The nudix hydrolases represent a class of proteins identified in many species including archaea, bacteria and eukarya (Xu *et al.*, 2001) that are known to catabolize substrates that are nucleoside diphosphates linked to a moiety X (Bessman *et al.*, 1996). They can be subdivided functionally according to their preference for dinucleoside polyphosphates, ADP-ribose, nucleotide sugars, NADH or various (deoxy)ribonucleoside triphosphates (Dunn *et al.*, 1999). nudix hydrolases are characterized by the presence of the highly conserved sequence motif (the nudix box) $GX_5EX_7UXEEXGU$ (where U is usually I, V or L and X can be any amino acid) that contributes to the catalytic and substrate-binding sites of the enzyme (Abeygunawardana *et al.*, 1995). The prototypical member of this family is the *Escherichia coli* MutT protein, which catalyzes the hydrolysis of nucleotide triphosphates with a preference for 8-oxo-dGTP, implying a role in DNA maintenance and repair.

Structures of functionally distinct nudix hydrolases from a variety of species have been solved (Abeygunawardana *et al.*, 1995; Lin *et al.*, 1997; Swarbrick *et al.*, 2000; Bonanno *et al.*, 2001; Durbecq *et al.*, 2001; Gabelli *et al.*, 2001; Bailey *et al.*, 2002; Fletcher *et al.*, 2002; Wang *et al.*, 2002). Some of these structures are monomers and some are dimers. All of them conserve a structural core of a central, mainly antiparallel, β -sheet and two helices (of which one is part of the loop-helix-loop containing the nudix box). The structural similarity of the core is extensive despite very low sequence identities; alignment of Ap₄A (diadenosine

5',5'''-P¹,P⁴-tetraphosphate) hydrolase sequences from animal and plant species reveals that even they have very low sequence identities (Swarbrick *et al.*, 2000; Bailey *et al.*, 2002; Fletcher *et al.*, 2002).

We are investigating the extremely radio-resistant bacterium *Deinococcus radiodurans* (Mattimore & Battista, 1996), which has been shown to encode at least 23 hypothetical nudix hydrolases, the largest number of any organism, which have diverse enzymatic activities and sizes (Makarova *et al.*, 2000; Xu *et al.*, 2001). As an initial step toward a structure–function analysis of the *D. radiodurans* nudix family, we report the expression, purification, crystallization and diffraction data collection for two members, DR1025 and DR0079. A multiple sequence alignment is shown in Fig. 1, comparing the *E. coli* MutT protein with DR1025 and DR0079 and illustrating the high sequence diversity outside the nudix box, even though a larger structural core is expected to be conserved for substrate binding and catalysis.

2. Materials and methods

2.1. Cloning, expression and purification

2.1.1. DR1025. The strain expressing DR1025 (pET24a expression vector in the Tuner cell background; Novagen) was obtained from Dr Maurice Bessman (Xu *et al.*, 2001). Cells were grown in Luria broth at 310 K with 30 µg ml⁻¹ kanamycin to an OD₆₀₀ of ~0.8. Protein expression was induced by addition of isopropyl- β -D-galactopyranoside (IPTG) to a concentration of 0.5 mM. Growth continued at




Figure 1
ClustalW multiple sequence alignment of *E. coli* MutT, DR1025 and DR0079. As shown (*, identities; ., similar; :, very similar), only the nudix box (bold) is conserved (8/23 identical). The rest of the sequences demonstrate the diversity of this protein family.

310 K for 3 h. Cells were harvested, rinsed and stored at 193 K. 3 g of cells was resuspended in 30 ml lysis buffer (50 mM Tris pH 7.5, 1 mM phenylmethylsulfonyl fluoride (PMSF), 10 $\mu\text{g ml}^{-1}$ DNase, 5 $\mu\text{g ml}^{-1}$ leupeptin, 10 $\mu\text{g ml}^{-1}$ antipain, 0.7 $\mu\text{g ml}^{-1}$ pepstatin, 0.3 $\mu\text{g ml}^{-1}$ chymostatin, 1 mM DTT) and lysed by sonication (pulsing) for 6 min. The cleared supernatant formed after centrifugation (12 000g, 20 min) was treated with streptomycin sulfate (1/2 volume of 10% solution) in order to precipitate nucleic acids. The resulting supernatant after centrifugation (12 000g, 20 min) was brought to 2.4 M ammonium sulfate to precipitate the proteins. The pellet formed after centrifugation (13 200g, 50 min) was resuspended in 25 mM Tris pH 7.5, 1 mM EDTA and dialyzed (8 kDa molecular-weight cutoff) against the same to remove salt. Anion-exchange chromatography (protein pI 4.76) was performed on a Pharmacia Hi-Trap Q HP column (5 ml) using the BioCAD Sprint system. Protein was eluted with a salt gradient from 0 to 1 M NaCl. DR1025-containing fractions (~0.4 M NaCl) were pooled, concentrated and desalted through buffer exchange using an Ultrafree centrifuge filter (5 kDa molecular-weight cutoff). SDS-PAGE showed the protein to be at least 98% pure and of proper molecular weight. Yield was approximately 14 mg pure protein from 1.5 g of cells (wet weight; 250 ml growth media) or 56 mg per litre of cells. Dynamic light-scattering experiments on a DynaPro99 (Protein Solutions) showed a radius of 2.52 nm and polydispersity of 42%. This is consistent with the protein being a monomer with a calculated molecular weight of 17 569 Da (159 amino acids).

2.1.2. DR0079. The DR0079 gene was amplified from the genomic DNA of *D. radiodurans* (ATCC) with the primers 5'-GACGGTCTCCCATGGGGGGCGTG-

AGC and 5'-GACGAGGAGGAATTCT-TATTCTCTCCTCCCTGTAGCAGC using standard PCR methods. The resulting PCR product was digested with *NcoI* and *EcoRI* and cloned into the pET30b expression vector (Novagen). DNA sequencing indicated that the sequence of the clone was identical to the reported sequence. The plasmid containing DR0079 was transformed into *E. coli* strain BL21(DE3) (Novagen) and cells were grown at 310 K to an OD₆₀₀ of ~0.8 in 750 ml Luria broth supplemented with 34 $\mu\text{g ml}^{-1}$ kanomycin. Protein expression was induced by addition of IPTG to a concentration of 1.0 mM. Following addition of 50 μl of 0.5 M zinc acetate, the cells were shaken for 4 h at 301 K, harvested (OD₆₀₀ of ~1.4) and frozen at 193 K. Thawed cells were resuspended in 40 ml lysis buffer (0.3 M NaCl, 50 mM potassium phosphate, 10 mM imidazole pH 7.5) and brought to 0.2 μM in PMSF prior to three passes through a French press (SLM Instruments). Following 1 min of sonication, the cell debris was spun at 24 000g for 45 min. The supernatant was loaded onto a 20 ml Ni-NTA affinity column (Qiagen) and washed stepwise with buffer (0.3 M NaCl, 50 mM potassium phosphate pH 8.1) containing increasing concentrations of imidazole (10–500 mM). The DR0079 fraction (~100 mg, 250 mM imidazole) was dialyzed (8 kDa molecular-weight cutoff) at 277 K overnight (50 mM NaCl, 20 mM Tris-HCl pH 7.6). Following volume reduction to ~3.0 ml (Amicon Centriprep-10), the N-terminal polyhistidine tag was removed at room temperature with 25 μl enterokinase (Novagen) and 6 μl of 1.0 M CaCl₂. After overnight incubation, an aliquot of the reaction mixture was loaded onto a Bio-Rad Econo-Pak High-Q cartridge (5 ml) using the BioCAD Sprint System (protein pI 4.85) to monitor the progress of the reaction. The column was washed (50 mM Tris-HCl,

50 mM Tris-bis propane pH 7.5) and eluted with a linear gradient from 0 to 150 mM NaCl over 20 column volumes. Under these conditions, cut DR0079 eluted with 55 mM NaCl and its fusion protein eluted with 90 mM NaCl. If the reaction was not over 90% complete, additional enterokinase was added.

Approximately 40 mg protein fractions were then loaded onto the High-Q cartridge. The peak containing DR0079 was collected, pooled and the volume reduced to ~2 ml. DR0079 was further purified on a Pharmacia Superdex75 HiLoad column that simultaneously exchanged it into crystallization buffer (20 mM Tris-HCl, 100 mM KCl, 1 mM DTT pH 7.1). Using a flow rate of 1.5 ml min⁻¹, approximately 18 mg of DR0079 eluted at 52 min, a retention time characteristic of a protein of size consistent with the calculated monomeric molecular weight of 19 283 Da (171 amino acids). The wide chemical shift dispersion and well defined cross peaks in the ¹⁵N/¹H HSQC NMR spectrum also suggest DR0079 to be a monomer in solution. SDS-PAGE showed the protein to be greater than 98% pure.

2.2. Crystallization and soaking

2.2.1. DR1025. Vapor-diffusion crystallization trials using hanging drops were set up using screens from Hampton Research. Large single crystals formed at room temperature after 1–2 d upon mixing equal volumes of protein (15 mg ml⁻¹ in 25 mM Tris pH 7.5, 1 mM EDTA) and reservoir (100 mM sodium acetate pH 4.5, 2 M sodium formate). The protein formed bipyramidal crystals of 200–500 μm in the longest dimension. A large single crystal is shown in Fig. 2(a). Crystals were prepared for freezing in liquid nitrogen by successively soaking for 5–10 min in reservoir solutions containing 10, 20 and 30% glycerol. Samarium-derivative crystals were prepared by soaking in 250 mM Sm(NO₃)₃, 0.74 M sodium acetate pH 4.5, 1.4 M sodium formate, 30% glycerol for 10 min or in 0.5 mM SmCl₃, 0.74 M sodium acetate pH 4.5, 1.4 M sodium formate for 24 h.

2.2.2. DR0079. Crystallization screening of DR0079 was performed at room temperature using the vapour-diffusion hanging-drop method and screens from Hampton Research. 2 μl of DR0079 (29 mg ml⁻¹) in crystallization buffer was mixed with 2 μl reservoir (30% PEG 4000, 0.1 M sodium citrate pH 5.6, 0.2 M ammonium acetate or 30% PEG 4000, 0.1 M Tris-HCl pH 8.5, 0.2 M lithium sulfate). Crystals formed as 'coffin-shaped' plates after

approximately four weeks and are shown in Fig. 2(b). Crystals were frozen in liquid nitrogen in the absence of additional cryoprotectant.

2.3. Diffraction data collection and reduction

All diffraction data were collected at the Macromolecular Crystallography Facility of the Advanced Light Source (ALS) synchrotron at Lawrence Berkeley National Laboratory. Heavy-atom incorporation into the DR1025 crystals was verified by fluorescence scanning. Images were integrated and scaled using the *HKL* and *HKL2000* packages of data-reduction software (Otwinowski, 1993). Crystal lattices and space groups for both proteins were determined by autoindexing and examination of systematic absences in the reduced data.

3. Results

3.1. Data collection for DR1025

Data from a native crystal were initially recorded to 1.6 Å resolution on ALS beamline 5.0.1. Subsequently, data were collected from a crystal that had been soaked for 24 h in SmCl₃ but did not show incorporation of Sm by fluorescence scans. This crystal diffracted to beyond 1.4 Å on ALS beamline 5.0.2. This data set is summarized in Table 1 and will be referred to as the 'native' data set. Assuming a

Table 1

Summary of crystallographic data.

Values in parentheses refer to the highest resolution shell (1.39–1.44 Å for DR1025 native; 1.80–1.84 Å for DR1025 Sm; 1.90–1.97 Å for DR0079).

| | DR1025 native | DR1025 Sm | DR0079 |
|---------------------------------|--|--|--|
| Wavelength (Å) | 1.0 | 1.2 | 1.0 |
| Crystal system | Tetragonal | Tetragonal | Orthorhombic |
| Space group | <i>P</i> 4 ₁ 2 ₁ 2/ <i>P</i> 4 ₂ 2 ₁ 2 | <i>P</i> 4 ₁ 2 ₁ 2/ <i>P</i> 4 ₂ 2 ₁ 2 | <i>C</i> 22 ₂ |
| Unit-cell parameters (Å) | <i>a</i> = <i>b</i> = 53.2, <i>c</i> = 122.6 | <i>a</i> = <i>b</i> = 53.2, <i>c</i> = 122.0 | <i>a</i> = 34.1, <i>b</i> = 157.2, <i>c</i> = 126.5 |
| Resolution (Å) | 1.39 | 1.80 | 1.90 |
| Total No. of observations | 879484 | 453884 | 642643 |
| Total No. of unique reflections | 35874 | 30510† | 26124 |
| Coverage completeness (%) | 98.4 (89.5) | 98.3 (98.2) | 95.4 (77.8) |
| <i>R</i> _{sym} ‡ (%) | 5.8 (34.4) | 4.8 (9.6) | 8.3 (36.7) |
| Average <i>I</i> /σ(<i>I</i>) | 56.1 (4.4) | 43.2 (18.9) | 22.7 (2.2) |

† Anomalous reflections not merged. ‡ $R_{\text{sym}} = \sum_{hkl} \sum_i |I_i - \langle I \rangle| / \sum_i I_i$, where I_i is the intensity of the *i*th measurement of reflection *hkl* and $\langle I \rangle$ is the mean intensity of reflection *hkl*.

monomer in the asymmetric unit, the estimated solvent content is 50.2% and the Matthews coefficient (*V*_M) is calculated to be 2.5 Å³ Da^{−1}.

Diffraction data were also collected from a crystal soaked in Sm(NO₃)₃ (see above). This crystal diffracted to 1.80 Å resolution with an *R*_{sym} of 4.8%, as shown in Table 1 (DR1025 Sm). Friedel pairs were collected by the inverse-beam procedure and reduced for SAD phasing. The presence of samarium was indicated by a fluorescence scan. The average anomalous difference for this data set was 5.0 and 7.3% (on *F*_{obs}) for the highest resolution range. A Harker section from the anomalous difference Patterson is shown in Fig. 3, illustrating several strong peaks from samarium sites.

3.2. Data collection for DR0079

Data from a native crystal were measured to 1.9 Å resolution on ALS beamline 5.0.2. The space group, unit-cell parameters and data-reduction statistics are given in Table 1. The diffraction indicated increased mosaicity of the crystals as data collection progressed. Assuming two protein molecules in the asymmetric unit, the estimated solvent content is 44.0% and the Matthews coefficient (*V*_M) is calculated to be 2.2 Å³ Da^{−1}, similar to that of DR1025.

4. Conclusions

In *D. radiodurans*, the nudix hydrolase family may contain up to 24 proteins, the most known for any organism. In light of the extreme radioresistance of this

bacterium and the activity of the prototypical nudix protein MutT in DNA repair, we are seeking to characterize the structure–function relationships of proteins of this family.

High-resolution native diffraction data were collected on two nudix family proteins from *D. radiodurans*, DR0079 and DR1025. Anomalous diffraction data were collected on a potential derivative of DR1025.

Dynamic light scattering (DR1025) and gel filtration (DR0079) suggest that both proteins are monomers in solution, even though DR0079 has two molecules per asymmetric unit. Previous structural studies have found nudix hydrolases both as monomers with one or two molecules per asymmetric unit (Swarbrick *et al.*, 2000; Bailey *et al.*, 2002) and as standard (Wang *et al.*, 2002) or domain-swapped dimers (Gabelli *et al.*, 2001). Solution-activity screens suggest that DR1025 may have activity towards dGTP and Ap₄A substrates, while the specificity of DR0079 is not apparent (Xu *et al.*, 2001). The three-dimensional structures of these proteins and their substrate complexes can be used to further characterize the functions of these proteins and the evolutionary relationships of this diverse family, as well as their role in the maintenance of cellular and genomic integrity. The data collected from the crystals described here is the initial step toward these goals.

This work was funded by a grant from the US Department of Energy, Office of Biological Energy Research, Contract No. DE-AC03-76SF00098. Diffraction data were collected at beamline 5.0.2 of the Macromolecular Crystallography Facility at the Advanced Light Source, Lawrence Berkeley




Figure 2
(a) Tetragonal crystal of DR1025. The longest dimension is approximately 500 μm. (b) Orthorhombic crystals of DR0079.




Figure 3
Harker section of anomalous difference Patterson for DR1025 samarium derivative. The *xz* plane is shown with the vertical axis *x* and the horizontal axis *z* in fractional coordinates from 0.0 to 0.5 for both axes. The three largest peaks of 21.8σ (0.5, 0.5, 0.0171), 11.1σ (0.1105, 0.5, 0.2194) and 10.6σ (0.0899, 0.5, 0.2794) can be seen.

crystallization papers

National Laboratory. Clones of DR1025 were constructed by Dr W. Xu and provided by Professor Maurice Bessman, Johns Hopkins University. We are grateful to Drs Emma Hill and Wasantha Ranatunga for assistance in preparation of this manuscript and to Ramona Pufan and Jana Mooster for their help in initial crystallization and derivatization of DR1025.

References

- Abeygunawardana, C., Weber, D. J., Gittis, A. G., Frick, D. N., Lin, J., Miller, A. F., Bessman, M. J. & Mildvan, A. S. (1995). *Biochemistry*, **34**, 14997–15005.
- Bailey, S., Sedelnikova, S. E., Blackburn, G. M., Abdelghany, H. M., Baker, P. J., McLennan, A. G. & Rafferty, J. B. (2002). *Structure*, **10**, 589–600.
- Bessman, M. J., Frick, N. & O'Handley, S. F. (1996). *J. Biol. Chem.* **271**, 25059–25062.
- Bonanno, J. B., Edo, C., Eswar, N., Pieper, U., Romanowski, M. J., Ilyin, V., Gerchman, S. E., Kycia, H., Studier, F. W., Sali, A. & Burley, S. K. (2001). *Proc. Natl Acad. Sci. USA*, **98**, 12896–12901.
- Dunn, C. A., O'Handley, S. F., Frick, D. N. & Bessman, M. J. (1999). *J. Biol. Chem.* **274**, 32318–32324.
- Durbecq, V., Sainz, G., Oudjama, Y., Clantin, B., Bompard-Gilles, C., Tricot, C., Caillet, J., Stalon, V., Droogmans, L. & Villeret, V. (2001). *EMBO J.* **20**, 1530–1537.
- Fletcher, J. I., Swarbrick, J. D., Maksel, D., Gayler, K. R. & Gooley, P. R. (2002). *Structure*, **10**, 205–213.
- Gabelli, S. B., Bianchet, M. A., Bessman, M. J. & Amzel, L. M. (2001). *Nature Struct. Biol.* **8**, 467–472.
- Lin, J., Abeygunawardana, C., Frick, D. N., Bessman, M. J. & Mildvan, A. S. (1997). *Biochemistry*, **36**, 1199–1211.
- Makarov, K. S., Aravind, L., Daly, M. J. & Koonin, E. V. (2000). *Genetica*, **108**, 25–34.
- Mattimore, V. & Battista, J. R. (1996). *J. Bacteriol.* **178**, 633–637.
- Otwinowski, Z. (1993). *Proceedings of the CCP4 Study Weekend. Data Collection and Processing*, edited by L. Sawyer, N. Isaacs & S. Bailey, pp. 56–62. Warrington: Daresbury Laboratory.
- Swarbrick, J. D., Bashtannyk, T., Maksel, D., Zhang, X. R., Blackburn, G. M., Gayler, K. R. & Gooley, P. R. (2000). *J. Mol. Biol.* **302**, 1165–1177.
- Wang, S., Mura, C., Sawaya, M. R., Cascio, D. & Eisenberg, D. (2002). *Acta Cryst. D* **58**, 571–578.
- Xu, W., Shen, J., Dunn, C. A., Desai, S. & Bessman, M. J. (2001). *Mol. Microbiol.* **39**, 286–290.

INTERNATIONAL UNION OF CRYSTALLOGRAPHY

Electronic Proof Instructions

These proofs should be returned **within 14 days of February 12 2003**. After this period, the Editors reserve the right to publish articles with only the Managing Editor's corrections.

Please

- (1) Read these proofs and assess if any corrections are necessary.
- (2) Check that any technical editing queries have been answered.
- (3) Return any corrections **immediately**

(a) by e-mail to

`lj@iucr.org`

giving a full description of the corrections in plain text and indicating the page, column and line numbers where appropriate.

(b) by fax to

+44 1244 314888

If the corrections are too complicated to send by e-mail please mark your corrections on the proofs as indicated overleaf. It is recommended that authors check that the fax transmission has been successful.

- (4) Authors are not required to return proofs by mail. However, if you wish to do this please send them to

Mr P. Strickland
Managing Editor,
International Union of Crystallography,
5 Abbey Square,
Chester CH1 2HU,
England

Telephone: +44 1244 342878

Substantial alterations, apart from occasioning delay in publication, are much more expensive than many authors would suppose. Authors may therefore be required to pay for any major alterations from their original copy, and it may sometimes be necessary to disallow such changes. Where alterations are unavoidable every effort should be made to substitute words or phrases equal in length to those deleted.

Please note that in order to save postal expenses and clerical work, the typescript, drawings and photographs of articles are normally destroyed after publication. Drawings and photographs will only be returned to authors if their return is specifically requested.

Authors will be informed by e-mail when their paper is published and may then download an electronic offprint of their paper from the author services page of **Crystallography Journals Online** (<http://journals.iucr.org>)

Printed offprints may be purchased using the attached form which should be returned as soon as possible. It will not be possible to supply offprints for orders received after the journal is printed.

Proof Corrections

If returning your corrections by fax (or mail) please use the marking system below. Please ensure that you use black ink (not pencil) and that you do not make corrections too near the edge of the page as these may be lost in transmission.

Please do not make corrections to the pdf file electronically and please do not return the pdf file.

The recommended system of marking corrections is as follows.

- (1) Each place in the text where a correction is needed should be indicated either by crossing out the characters to be corrected or by an insert mark (\wedge) if an insertion is needed.
- (2) The correct character, word, instruction or insertion should be shown in the margin at one end of the line and should be terminated with a long slash mark (\swarrow).
- (3) The corrections in the margins (both left and right margins may be used) should be arranged in the same sequence as they are required in the line of text.
- (4) Characters to be printed in the superior position, such as superscripts, apostrophes and quotation marks, are to be so identified by the symbol \vee under them, e.g. $\vee\text{V}$ or $\vee\text{V}$. Characters to be printed in the inferior position, such as subscripts, are to be identified by the symbol \wedge over them, e.g. $\wedge\text{A}$.
- (5) Instructions written in the margins should be encircled. This indicates that this text is not to be printed. This does not apply to the full stop (period), the colon and the solidus (slash mark), which are circled for clarity.
- (6) Some instructions are needed so frequently that it is convenient to indicate them simply with special symbols or abbreviations. The symbols are recognized as instructions without being circled. Some frequently used instructions are given in the following list.

| Alteration | Marginal mark | Mark in text |
|--|-----------------------|---|
| Insert or substitute: | | |
| Space | #/ | \wedge (for an insertion) or delete material to be replaced |
| Full stop (period) | ⊙/ | |
| Colon | ⊖/ | |
| Solidus (slash mark) | ⊘/ | |
| Hyphen | - | |
| Rule | ⊗/ | |
| Chemical bond | ⊕/ | |
| Superior (e.g. superscript 2 or apostrophe) | $\vee/$ or $\vee/$ | |
| Inferior (e.g. subscript 2) | $\wedge\text{A}$ | |
| Change to: | | |
| Capitals | <u>Cap</u> | \equiv \equiv \equiv } under characters |
| Small capitals | <u>Sc</u> | |
| Italic type | <u>Ita</u> | |
| Bold type | <u>Bold</u> | |
| Lower case letters | <u>Lc</u> | } Circle characters |
| Roman type | <u>Rom</u> | |
| Delete | ? | Cross out unwanted material |
| Delete and close up | $\overline{\text{?}}$ | Cross out unwanted material and surround with $\overline{\text{?}}$ |
| Close up | \sim | \sim around space to be closed up |
| Invert type | ⊖ | Circle inverted characters |
| Transpose | ⊗ | \sqcap between letters or words |
| Faulty setting (e.g. broken type) | X | Circle defective characters |
| Leave as printed | <u>stet</u> | under material to be left |
| New paragraph | <u>¶</u> | \sqcap before first word of new paragraph |
| No new paragraph or line | <u>run on</u> | \hookrightarrow between lines |

Please note

- (a) No correction should be made in the text without an accompanying mark in the margin, or the correction may be easily missed.
- (b) Complicated corrections can be explained in a covering letter.

PAID OFFPRINT ORDER FORM

Return within 14 days of February 12 2003

YOU WILL AUTOMATICALLY BE SENT DETAILS OF HOW TO DOWNLOAD
AN ELECTRONIC OFFPRINT OF YOUR PAPER, FREE OF CHARGE.
PRINTED OFFPRINTS MAY BE PURCHASED USING THIS FORM.

Please keep a copy of this order and send the original to:

INTERNATIONAL UNION OF CRYSTALLOGRAPHY
5 Abbey Square
Chester CH1 2HU, England

VAT No. GB 161 9034 76

Article No.: D030267-PU0091

Author(s) Elizabeth L. Holbrook, Ursula Schulze-Gahmen, Garry W. Buchko, Shuisong Ni, Michael A. Kennedy, Stephen R. Holbrook,

Title of article Purification, crystallization and preliminary X-ray analysis of two nudix hydrolases from *Deinococcus radiodurans*

E-mail address (for electronic offprints) sholbrook@lbl.gov

I wish to order paid offprints

These offprints should be sent to

Name Professor Stephen R. Holbrook

Address Structural Biology Dept, Physical Sciences Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Bldg 64-123, Berkeley, CA 94720, USA

If the above address or e-mail address is not correct, please indicate an alternative:

| |
|--|
| |
|--|

Date

Signature

☐ A cheque for payable to **INTERNATIONAL UNION OF CRYSTALLOGRAPHY** is enclosed

☐ An official purchase order made out to **INTERNATIONAL UNION OF CRYSTALLOGRAPHY** ☐ is enclosed ☐ will follow

Purchase order No.

☐ Please invoice me

OFFPRINTS

An electronic offprint is supplied free of charge.

Printed offprints without limit of number may be purchased at the prices given in the table below. The requirements of all joint authors, if any, and of their laboratories should be included in a single order, specifically ordered on the form overleaf, which must be accompanied by a remittance in **United States dollars** payable to **INTERNATIONAL UNION OF CRYSTALLOGRAPHY** unless an official purchase order made out to **INTERNATIONAL UNION OF CRYSTALLOGRAPHY** is sent. All orders must be submitted promptly; it will not be possible to supply offprints for orders received after the journal is printed.

Please note that normally offprints are sent about one month after publication of the article, and normally by surface mail. If offprints are wanted by air mail additional costs will be invoiced.

Orders should be returned by fax to:

INTERNATIONAL UNION OF CRYSTALLOGRAPHY
5 Abbey Square
Chester CH1 2HU, England
Fax: +44 1244 314888

PRICE LIST OF OFFPRINTS

Prices are given in **United States dollars** and include postage.

Offprints with covers are only available by special order. Please contact the Managing Editor (at the address given above) for a quotation.

| Number of offprints required | Size of paper (in printed pages) | | | | |
|------------------------------|----------------------------------|-----|-----|------|----------------|
| | 1–2 | 3–4 | 5–8 | 9–16 | Additional 8's |
| 50 | 42 | 60 | 83 | 126 | 56 |
| 100 | 61 | 85 | 118 | 180 | 73 |
| 150 | 80 | 110 | 154 | 234 | 92 |
| 200 | 99 | 136 | 190 | 289 | 114 |
| Additional 50's | 19 | 25 | 36 | 55 | 18 |

ENQUIRIES

Enquiries concerning offprints should be sent to support@iucr.org.